



Primary prevention of sexually transmitted disease: applying the ABC strategy

S J Genuis and S K Genuis

Postgrad. Med. J. 2005;81;299-301
doi:10.1136/pgmj.2004.026039

Updated information and services can be found at:
<http://pmj.bmjournals.com/cgi/content/full/81/955/299>

These include:

Rapid responses

You can respond to this article at:
<http://pmj.bmjournals.com/cgi/eletter-submit/81/955/299>

Email alerting service

Receive free email alerts when new articles cite this article - sign up in the box at the top right corner of the article

Topic collections

Articles on similar topics can be found in the following collections

[HIV Infection/AIDS](#) (1021 articles)
[Prevention and health promotion](#) (603 articles)
[Sexually Transmitted Infections](#) (1055 articles)

Notes

To order reprints of this article go to:
<http://www.bmjournals.com/cgi/reprintform>

To subscribe to *Postgraduate Medical Journal* go to:
<http://www.bmjournals.com/subscriptions/>

REVIEW

Primary prevention of sexually transmitted disease: applying the ABC strategy

S J Genuis, S K Genuis

Postgrad Med J 2005;81:299–301. doi: 10.1136/pgmj.2004.026039

Escalating rates of sexually transmitted disease (STD) in many areas of the world necessitate a re-evaluation of current public health STD preventive programmes. Pervasive long term sequelae for many STD afflicted people and the emerging threat, caused by the HIV/AIDS pandemic, to some national infrastructures, suggests that ongoing initiatives focusing primarily on risk reduction through barrier protection have not met their desired objective. Recent strategies to promote non-coital sexual involvement as a means of achieving STD reduction fail to address the transmission of infection that may occur through alternative non-intercourse sexual activities. The demonstrated success of the innovative, comprehensive ABC strategy shows that while risk reduction and treatment of existing infection remain important, the promotion of optimal health may be achieved more effectively through broad based comprehensive and adaptable programmes that include an emphasis on risk avoidance through delayed sexual debut and partner reduction.

programmes principally focus on the reduction of risk as a means of stemming the tide of these diseases. The promotion of “safe” or “safer” sex practices and, more recently, of non-coital sexual activity has resulted in a focus on risk reduction even in population groups where an emphasis on risk elimination might be appropriate.

A focus on condom use, as a means of risk reduction, has been the basis of most behavioural STD interventions for many years. The practical effectiveness of this strategy is, however, hindered by three important factors. Firstly, condoms provide limited protection against “SS” (skin to skin or skin to sore) transmission of STDs such as HPV, HSV, syphilis, or chancroid as these barrier devices do not cover all exposed areas and thus direct skin contact with pathogens throughout the external genital area compromises the protection offered by the condom.¹ Secondly, rates of mechanical failure and user error with condoms, particularly in young people, are significant.^{13–14} And finally, issues of compliance seriously impair the effectiveness of protection against discharge related infections such as HIV, chlamydia, and gonorrhoea.¹ Research shows that exposure to condom education and increased awareness of risk does not result in safer sex choices when adolescents are sexually aroused, and only a minority of people engaging in risky sexual behaviour use condoms consistently.^{15–19} This latter point has serious implications as “irregular use of condoms provides no protection against transmission of HIV and STD.”²⁰

Despite recent reports of decreased STD rates as a result of widespread condom use by commercial sex workers (CSW),^{21–22} published results suggest that partner reduction played a pivotal part in STD decline. Mass advertising campaigns and widespread HIV education in Thailand as well as Ethiopia, Cambodia, and other countries resulted in pronounced reduction of casual sex and liaisons with CSW.^{6–23–25} Infection patterns suggest the central role of partner reduction: after the “100% condom program” for Thailand’s CSW, for example, there was a pronounced decrease not only in discharge related STDs, but also in “SS” diseases such as syphilis and chancroid,^{21–22} infections that are commonly transmitted despite condom use. Furthermore, in an article comparing HIV seroprevalence among CSW who started the sex trade before and after the “100% condom program”, it was found that HIV seroprevalence

Despite international efforts to reduce the incidence of sexually transmitted disease (STD), rising infection rates^{1–4} highlight the importance of exploring and evaluating the efficacy of STD prevention strategies. While the presentation of factual information about barrier protection and infection management has been the mainstay of prevention programmes worldwide, an overview of current STD risk reduction strategies and an exploration of programmes that focus on primary prevention suggests that the ABC approach to sexual behaviour change may be effectively used to develop balanced, population specific STD prevention programmes.^{5–6} This approach, which advocates delayed sexual debut for adolescents (A, abstinence), partner reduction (B, be faithful), and factual information regarding condom use (C, condoms)^{5–6} has been successfully pioneered in Uganda, a country previously experiencing significant STD rates.⁷ Perhaps it is time to consider the potential application of this approach in the educational programmes of other nations.^{6–8–11}

CURRENT RISK REDUCTION STRATEGIES

Rising STD rates have led to a situation where certain sexually transmitted infections are regarded as “an inevitable consequence of sexual activity”¹² and current STD prevention

Abbreviations: STD, sexually transmitted disease; CSW, commercial sex worker

See end of article for authors’ affiliations

Correspondence to:
Dr S J Genuis, 2935-66
Street, Edmonton, Alberta,
Canada T6K 4C1;
sgenuis@ualberta.ca

Submitted 24 June 2004
Accepted 5 August 2004

was higher among CSW who started the sex trade after the implementation of the mandatory condom programme,²⁵ thus suggesting ongoing vulnerability for CSW and clients despite legislated condom use. In addition, it has been found that many Thai citizens, particularly adolescents, are still not consistently using condoms.^{16 26 27} These findings imply that less commercial sex and partner reduction played an important part in the falling STD rates in these nations^{6 23 25} and supports the hypothesis that a comprehensive educational strategy that incorporates risk avoidance through partner reduction is likely to have a greater impact than strategies with unilateral focus on risk reduction through condoms.

While factual information regarding barrier protection is an important component of STD reduction strategies, it has not met the desired objective of diminishing the global STD pandemic. As a result of concerns about escalating STDs in the adolescent population (the World Health Organisation estimates that two thirds of STDs worldwide occur in teenagers and young adults²⁸) and the increasing recognition that, from an STD perspective, early sexual intercourse among youth may be a health hazard, some educational initiatives have sought to promote non-coital sexual activity as a means of STD risk reduction. This strategy, however, has limited potential for STD prevention: non-intercourse sexual involvement may entail serious risk of STD acquisition.¹⁴ Oral sex, for example, may be a means of transmitting both "SS" and discharge related STDs, and genital frottage facilitates transmission of "SS" pathogens.^{14 29 30} In addition, there is no evidence to support the hypothesis that young people participating in highly stimulating, non-coital sexual activity will not participate in at least intermittent intercourse.

With the limited success of programmes focusing primarily on risk reduction, attention to the ABC strategy for sexual behaviour change, particularly the often neglected A and B components, is warranted.^{6 9}

PRIMARY PREVENTION: ADDRESSING UNDERLYING BEHAVIOURS

There has been general consensus in the public health community that HIV discordant couples and those engaging in higher risk sexual activities, including multiple sexual partners or "serial monogamy", should be encouraged to use condoms to reduce the transmission of discharge related STDs; however, focus on this risk reduction strategy has sometimes occurred without a complementary focus on other forms of behavioural interventions. The fact that early initiation of sexual intercourse in adolescents is associated with a higher number of lifetime sexual partners³¹⁻³⁴ and a consequent higher risk of contracting STDs, makes the promotion of delayed sexual debut in this population a first and critical step in partner reduction and the primary prevention of sexually transmitted infections.⁹ From a STD perspective, sexual encounters include present partners as well as past partners and all of their partner's partners³⁵; therefore, partner reduction must be a prerequisite focus in the primary prevention of STDs. The results of a prevention programme that focused on the both primary prevention and risk reduction through the use of an ABC approach to sexual behaviour change can be seen in the outcomes of a national programme started by government and health officials in Uganda.⁵⁸

To tackle the growing HIV crisis, an aggressive media campaign was instituted in Uganda over the past decade with a strong emphasis on "behaviour change" in the general population as well as an emphasis on empowering women to make healthy sexual choices.⁵ Reported changes occurring in association with this programme include the following; rates of 13 to 16 year olds involved in sexual activity in one district

of Uganda declined from nearly 60% in 1994 to less than 5% by 2001⁵; fewer than 10% of unmarried Ugandan women in all age groups reported multiple partners, in noticeable contrast with 20%-65% of women in other African nations such as Kenya and Malawi⁵; and, reported rates of barrier protection use with non-regular partners increased. During about the same time period, national HIV prevalence declined from reported rates as high as 30% in 1992⁷ to an estimated 5% in 2001.⁵ The potential public health implications of replicating the Ugandan results in other nations are apparent.

It has been reported that "behavior change, as distinct from condom adoption"²⁴ was the principal factor responsible for pronounced reduction in HIV rates in Uganda.^{5 6 8 24} Although barrier protection with irregular partners increased in the past half decade, ever use of condoms in Uganda remained low: only 16% of women in 2000 reported that they had ever had sex with men who were using condoms,⁵ and a study of the general population in one district found that only 4.4% reported consistent condom use.²⁰ In other African nations, such as Zimbabwe and Botswana, where STD prevention strategies have principally focused on the promotion and distribution of condoms and where, compared with Uganda, there are higher rates of condom sales and reported use, HIV prevalence is among the highest in the world.^{5 36-38} There is an unprecedented 55.6% HIV prevalence among pregnant women aged 25-29 in urban Botswana and infection rates in Zimbabwe suggest that by 2020 there will be a 30% AIDS related workforce loss.³⁹ These figures contrast starkly with Ugandan figures and suggest that the ABC approach may provide an adaptable framework for balanced, population specific STD prevention programmes in other nations.⁸

CONCLUSION

The worldwide STD crisis, a public health challenge that disregards socioeconomic barriers and national boundaries, requires broad based interventions that allow for personal, cultural, and epidemiological differences. Given the apparent success of prevention strategies that address primary sexual behaviour, increased consideration and resources should be allocated to comprehensive ABC STD prevention initiatives that include the promotion of risk avoidance through delayed sexual debut and partner reduction, as well as the provision of factual information about risk reduction through barrier protection.

Authors' affiliations

S J Genuis, Department of Obstetrics and Gynaecology, University of Alberta, Canada

S K Genuis, Research Librarian, Alberta, Canada

Funding: none.

Conflicts of interest: none.

Dr Stephen J Genuis served as Co-Director of Medical Services at a hospital in Cameroon. Shelagh K Genuis writes on topics related to information and health. SJG and SKG recently coauthored *Teen Sex: Reality Check*. Both authors were involved in the preparation of this paper.

REFERENCES

- Workowski KA**, Levine WC. Sexually transmitted diseases treatment guidelines 2002. Centers for Disease Control and Prevention. *MMWR Recomm Rep* 2002;**51**:1-78.
- Fenton K**, Giesecke J, Hamers FF. Europe-wide surveillance for sexually transmitted infections: a timely and appropriate intervention. *Euro Surveill* 2001;**6**:69-70.
- Weiss RA**, Adler MW, Rowland-Jones SL. The changing face of HIV and AIDS: preface. *Br Med Bull* 2001;**58**:1-2.
- Workowski KA**, Levine WC, Wasserheit JN. US Centers for Disease Control and Prevention guidelines for the treatment of sexually transmitted diseases: an opportunity to unify clinical and public health practice. *Ann Intern Med* 2002;**137**:255-62.

- 5 Hogle J, Green EC, Nantulya V, *et al.* *What happened in Uganda? Declining HIV prevalence, behavior change, and the national response.* Washington DC: US Agency for International Development, Office of HIV/AIDS, Bureau of Global Health, 2002.
- 6 Shelton AJ, Halperin DT, Nantulya V, *et al.* Partner reduction is crucial for balanced "ABC" approach to HIV prevention. *BMJ* 2004;**328**:891-4.
- 7 Anon. Uganda tackles AIDS from the very top down. *AIDS Alert* 1999;**14**(suppl 8):3-4.
- 8 Stoneburner RL, Low-Beer D. Population-level HIV declines and behavioral risk avoidance in Uganda. *Science* 2004;**304**:714-18.
- 9 Genuis SJ, Genuis SK. Adolescent behaviour should be priority. *BMJ* 2004;**328**:894.
- 10 Wilson D. Partner reduction and the prevention of HIV/AIDS. *BMJ* 2004;**328**:848-9.
- 11 Stammers TG. Abstinence under fire. *Postgrad Med J* 2003;**79**:365-6.
- 12 Collins S, Mazloomzadeh S, Winter H, *et al.* High incidence of cervical human papillomavirus infection in women during their first sexual relationship. *B J Obstet Gynaecol* 2002;**109**:96-8.
- 13 Warner L, Clay-Warner J, Boles J, *et al.* Assessing condom use practices. Implications for evaluating method and user effectiveness. *Sex Transm Dis* 1998;**25**:273-7.
- 14 Genuis SJ, Genuis SK. Orgasm without organisms: science or propaganda? *Clin Pediatr* 1996;**35**:10-17.
- 15 de Vincenzi I. A longitudinal study of human immunodeficiency virus transmission by heterosexual partners. European Study Group on Heterosexual Transmission of HIV. *N Engl J Med* 1994;**331**:341-6.
- 16 Lertpiriyasuwat C, Pliat T, Jenkins RA. A survey of sexual risk behavior for HIV infection in Nakhonsawan, Thailand, 2001. *AIDS* 2003;**17**:1969-76.
- 17 Weisman CS, Plichta S, Nathanson CA, *et al.* Consistency of condom use for disease prevention among adolescent users of oral contraceptives. *Fam Plann Perspect* 1991;**23**:71-4.
- 18 Hausser D, Michaud PA. Does a condom-promoting strategy (the Swiss STOP-AIDS campaign) modify sexual behavior among adolescents? *Pediatrics* 1994;**93**:580-5.
- 19 Bankole A, Darroch JE, Singh S. Determinants of trends in condom use in the United States, 1988-1995. *Fam Plann Perspect* 1999;**31**:264-71.
- 20 Ahmed S, Lutalo T, Wawer M, *et al.* HIV incidence and sexually transmitted disease prevalence associated with condom use: a population study in Rakai, Uganda. *AIDS* 2001;**15**:2171-9.
- 21 Rojanapithayakorn W, Hanenberg R. The 100% condom program in Thailand. *AIDS* 1996;**10**:1-7.
- 22 Hanenberg RS, Rojanapithayakorn W, Kunasol P, *et al.* Impact of Thailand's HIV-control programme as indicated by the decline of sexually transmitted diseases. *Lancet* 1994;**344**:243-5.
- 23 Pisani E, Garnett GP, Grassly NC, *et al.* Back to basics in HIV prevention: focus on exposure. *BMJ* 2003;**326**:1384-7.
- 24 Green EC, Conde A. Sexual partner reduction and HIV infection. *Sex Transm Infect* 2000;**76**:145.
- 25 Kilmarx PH, Palanuvej T, Limpakarnjanarat K, *et al.* Seroprevalence of HIV among female sex workers in Bangkok: evidence of ongoing infection risk after the "100% condom program" was implemented. *J Acquir Immune Defic Syndr* 1999;**21**:313-16.
- 26 Paz-Bailey G, Kilmarx PH, Supawitkul S, *et al.* Risk factors for sexually transmitted diseases in northern Thai adolescents: an audio-computer-assisted self-interview with noninvasive specimen collection. *Sex Transm Dis* 2003;**30**:320-6.
- 27 Thato S, Charron-Prochownik D, Dorn LD, *et al.* Predictors of condom use among adolescent Thai vocational students. *J Nurs Scholarsh* 2003;**35**:157-63.
- 28 Dehne KL, Riedner G. Sexually transmitted infections among adolescents: the need for adequate health services. *Reprod Health Matters* 2001;**17**:170-83.
- 29 Giovannelli L, Campisi G, Lama A, *et al.* Human papillomavirus DNA in oral mucosal lesions. *J Infect Dis* 2002;**185**:833-6.
- 30 Edwards S, Carne C. Oral sex and the transmission of viral STIs. *Sex Transm Infect* 1998;**74**:6-10.
- 31 Sonenstein FL, Pleck JH, Ku LC. Levels of sexual activity among adolescent males in the United States. *Fam Plann Perspect* 1991;**23**:162-7.
- 32 Greenberg J, Magder L, Aral S. Age at first coitus. A marker for risky sexual behavior in women. *Sex Transm Dis* 1992;**19**:331-4.
- 33 Wasserheit JN. Effect of changes in human ecology and behavior on patterns of sexually transmitted diseases, including human immunodeficiency virus infection. *Proc Natl Acad Sci USA* 1994;**91**:2430-5.
- 34 Santelli JS, Brener ND, Lowry R, *et al.* Multiple sexual partners among U.S. adolescents and young adults. *Fam Plann Perspect* 1998;**30**:271-5.
- 35 Genuis S, Genuis SK. *Teen sex: reality check.* Edmonton, Alberta: Winfield House Publishing, 2002.
- 36 Centres for Disease Control. The global HIV and AIDS epidemic, 2001. *MMWR Morb Mortal Wkly Rep* 2001;**50**:434-9.
- 37 Allen A. Sex change. *The New Republic* 2002;**226**:14-15.
- 38 Adetunji J, Meekers D. Consistency in condom use in the context of HIV/AIDS in Zimbabwe. *J Biosoc Sci* 2001;**33**:121-38.
- 39 Fact list. AIDS: focus on Africa. *Can Med Assoc J* 2002;**167**:529.